

IN THE CLAIMS:

Applicants, pursuant to 37 C.F.R. § 1.121, submit the following amendments to the claims:

1. (Currently amended) A diagnostic ~~or prognostic~~ assay for cancer, comprising:
  - (a) obtaining a tissue sample from a test tissue;
  - (b) performing a methylation assay on DNA from the tissue sample, wherein the methylation assay determines the methylation state of one or more coordinately methylated CpG dinucleotide sequences within SEQ ID NO:36; and
  - (c) determining a diagnosis ~~or prognosis~~ based, at least in part, upon the methylation state of the CpG dinucleotide within the DNA sequence, compared to that of control DNA, wherein the determined methylation state is hypermethylation, and wherein the cancer is breast cancer.
- 2.-3. (Cancelled)
4. (Currently amended) The diagnostic ~~or prognostic~~ assay of claim 1 wherein the methylation assay procedure is selected from the group consisting of MethyLight, MS-SNuPE, MSP, MCA, COBRA, and combinations thereof.
- 5.-6. (Cancelled)
7. (Previously presented) A kit useful for the detection of a methylated CpG-containing nucleic acid comprising a carrier means containing one or more containers comprising:
  - (a) a container containing a probe or primer consisting of at least 12 contiguous nucleotides of a sequence selected from the group consisting of SEQ ID NOS:36 and 37, and the bisulfite-converted sequences thereof; and
  - (b) additional standard methylation assay reagents, wherein the kit, based at least in part on the probe or primer, is suitable to determine the methylation status of one or more CpG dinucleotides within the sequence selected from the group consisting of SEQ ID NOS:36 and 37.
8. (Previously presented) The kit of claim 7, wherein the additional standard methylation assay reagents are standard reagents for performing a methylation assay from the group consisting of MethyLight, MS-SNuPE, MSP, MCA, COBRA, and combinations thereof.

9. (Cancelled)

10. (Previously presented) An isolated nucleic acid molecule consisting of a methylated or unmethylated polynucleotide sequence selected from the group consisting of sequences of SEQ ID NO:37 and the bisulfite-converted sequences thereof.

11. (Original) The nucleic acid of claim 10, wherein the nucleic acid is methylated.

12. (Original) The nucleic acid of claim 10, wherein the nucleic acid is unmethylated.

13. (Currently amended) A diagnostic ~~or prognostic~~ assay for cancer, comprising:

(a) obtaining a tissue sample from a test tissue;

(b) performing a methylation assay on DNA from the tissue sample, wherein the methylation assay determines the methylation state of a CpG dinucleotide within SEQ ID NO:37; and

(c) determining a diagnosis ~~or prognosis~~ based, at least in part, upon the methylation state of the CpG dinucleotide within the DNA sequence, compared to that of control DNA, wherein the determined methylation state is hypermethylation, and wherein the cancer is prostate, breast or colon cancer.

14. (Cancelled)

15. (Currently amended) The diagnostic ~~or prognostic~~ assay of claim 13, wherein the methylation assay procedure is selected from the group consisting of MethyLight, MS-SNuPE, MSP, MCA, COBRA, and combinations thereof.

16. (Currently amended) A diagnostic ~~or prognostic~~ assay for cancer, comprising:

(a) obtaining a tissue sample from a test tissue;

(b) performing a methylation assay on DNA from the tissue sample, wherein the methylation assay determines the methylation state of a CpG dinucleotide within SEQ ID NO:36; and

(c) determining a diagnosis ~~or prognosis~~ based, at least in part, upon the methylation state of the CpG dinucleotide, compared to that of control DNA, wherein hypermethylation of SEQ ID NO:36 is indicative of breast cancer.

17. (Currently amended) The diagnostic ~~or prognostic~~ assay of claim 16, wherein the methylation assay procedure is selected from the group consisting of MethyLight, MS-SNuPE,

MSP, MCA, COBRA, and combinations thereof.

18. (Currently amended) A diagnostic ~~or prognostic~~ assay for cancer, comprising:

(a) obtaining a tissue sample from a test tissue;

(b) performing a methylation assay on DNA from the tissue sample, wherein the methylation assay determines the methylation state of a CpG dinucleotide within SEQ ID NO:37; and

(c) determining a diagnosis ~~or prognosis~~ based, at least in part, upon the methylation state of the CpG dinucleotide within the DNA sequence, compared to that of control DNA, wherein hypermethylation of SEQ ID NO:37 is indicative of prostate, breast or colon cancer.

19. (Currently amended) The diagnostic ~~or prognostic~~ assay of claim 18, wherein the methylation assay procedure is selected from the group consisting of MethyLight, MS-SNuPE, MSP, MCA, COBRA, and combinations thereof.